

CORRECTED
VERSION*

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



RECEIVED

DEC 11 2000

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12Q 1/68 A1 (11) International Publication Number: WO 00/22167 (43) International Publication Date: 20 April 2000 (20.04.00)

(21) International Application Number: PCT/US99/23277

(22) International Filing Date: 6 October 1999 (06.10.99)

(30) Priority Data:
60/103,803 9 October 1998 (09.10.98) US

(71) Applicant (for all designated States except US): XGENE CORPORATION [US/US]; 863C Mitten Road, Burlington, CA 94010 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): HOFFFLER, Warren [US/US]; 2028 Howard Avenue, San Carlos, CA 94070 (US).

(74) Agents: WARD, Michael, R. et al.; Limbach & Limbach L.L.P., 2001 Ferry Building, San Francisco, CA 94111-4262 (US).

(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

With international search report.
Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: METHOD FOR DETERMINING TRANSCRIPTION FACTOR ACTIVITY AND ITS TECHNICAL USES

(57) Abstract

Certain transcription factors (enhancer binding proteins) significantly increase transcription rates from genes by nicking a single DNA strand in the vicinity of their DNA binding sites, thereby allowing RNA polymerase to gain access to the transcribed DNA strand by a process of "threading". DNA template nicking is a detectable and quantifiable byproduct indicative of transcriptional activation that can be used to design practical assays. These assays are used to determine which transcription factors (enhancer binding proteins) are actively catalyzing the transcription of a gene in any cell type, or in any cell in response to any drug or treatment. This group of transcription factors have a predictable molecular biological activity in addition to transcription activation, namely site-specific DNA strand cleavage.

RECEIVED
JAN 22 2001
TO ITD MAIL ROOM